



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE FUNCTIONAL EQUATION $g(x^2) = 2ax + [g(x)]^2$.

BY J. H. M. WEDDERBURN.

1. **Introduction.** The functional equation considered in this paper arose out of an extension of a problem in arrangements which occurs in the theory of linear algebras. In an algebra which is neither associative nor commutative, n factors may be associated in a number of different ways; thus for four factors a_1, a_2, a_3, a_4 we have five different associations in which the subscripts all occur in their natural order, namely

$$\begin{array}{lll} a_1 (a_2 \cdot a_3 a_4), & (a_1 a_2) (a_3 a_4), & (a_1 \cdot a_2 a_3) a_4, \\ a_1 (a_2 a_3 \cdot a_4), & & (a_1 a_2 \cdot a_3) a_4. \end{array}$$

The first problem then is to determine the number N_n of such *types* of association for n factors.

It is easily seen* that we can count the number of different types by taking first those in which the left-hand factor consists of one element and the right-hand one of $n-1$, then those in which the first has two elements and the second $n-2$, and so on. Hence

$$N_n = N_1 N_{n-1} + N_2 N_{n-2} + \cdots + N_{n-1} N_1.$$

If we set

$$f(x) = N_1 x + N_2 x^2 + N_3 x^3 + \cdots,$$

we have, since $N_1 = 1$,

$$f^2(x) = f(x) - x,$$

and therefore, since $f(x)$ vanishes for $x = 0$,

$$f(x) = \frac{1 - \sqrt{1 - 4x}}{2} = \sum_1^{\infty} \frac{(2n-2)!}{(n-1)! n!} x^n,$$

so that

$$N_n = \frac{(2n-2)!}{(n-1)! n!}.$$

* This solution (for an equivalent problem) is also given by P. Quarra, *Torino Atti* vol. 53, (1918) pp. 1044-1047. See also P. Franklin, *Question 2681*, *Amer. Math. Monthly* vol. 25 (1918), p. 118 and solution by C. F. Gummer, *ibid.*, vol. 26 (1919) pp. 127-128.

If we assume that multiplication is commutative, the problem of counting types is much more difficult as is seen from the fact that for $n = 4$

$$a_1 (a_2 \cdot a_3 a_4), \quad (a_1 a_2) (a_3 a_4)$$

are the only types since $a_1 \cdot a_2 a_3 = a_2 a_3 \cdot a_1$ are now of the same type. If n is odd, the method of counting used above is still valid and leads to

$$(1) \quad 2 N_n = N_1 N_{n-1} + N_2 N_{n-1} + \cdots + N_{n-1} N_1,$$

since the types corresponding to the term $N_{n-r} N_r$ are the same as those belonging to the term $N_r N_{n-r}$. If, however, n is even, say $n = 2k$, the number of types in which the number of elements in the left and right factors is $k = n/2$ is obviously not $N_k^2/2$ but $N_k(N_k + 1)/2$, so that we have in place of (1)

$$(2) \quad 2 N_n = N_1 N_{n-1} + N_2 N_{n-2} + \cdots + N_{n-1} N_1 + N_{\frac{n}{2}}.$$

If, on the analogy of the solution in the previous problem, we set

$$g(x) = 1 - \sum_1^{\infty} N_n x^n,$$

we readily derive from (1) and (2) that $g(x)$ satisfies the functional equation

$$g(x^2) = 2x + g^2(x).$$

This equation may be replaced by another of somewhat simpler form by setting

$$(3) \quad h(x) = g(x)/x^{\frac{1}{2}},$$

which gives

$$h(x^2) = 2 + h^2(x),$$

from which it is obvious that $g_s = h(x^{2^s})$ is a solution of the difference equation

$$g_{s+1} = 2 + g_s^2.$$

This suggests the consideration of the difference equation

$$(4) \quad \psi_{s+1} = a\psi_s^2 + b\psi_s + c$$

which may be written

$$(8') \quad a_1 = \alpha, \quad 2a_m = \sum_{r=1}^{m-1} a_r a_{m-r} + a_{m/2}, \quad (m = 2, 3, \dots)$$

if we agree to reckon $a_{m/2}$ as zero when $m/2$ is not an integer.

There are two exceptional values of α in which the solution (7) is trivial; firstly $\alpha = 0$, which gives $g(x) \equiv 1$, and secondly $\alpha = -1$, in which case all the a 's vanish except the first and

$$g(x) = 1 + x, \quad (\alpha = -1).$$

The case $\alpha = 0$ is excluded from further consideration unless specially mentioned and, of course, in the case $\alpha = -1$ the discussion in the remaining sections is trivial.

3. The convergence of the series for $g(x)$. If the term $a_{m/2}$ in (8') is suppressed, equation (5) becomes

$$(9) \quad g^2(x) = -2\alpha x + 1,$$

or

$$\begin{aligned} g(x) &= \sqrt{1 - 2\alpha x} = 1 - \alpha x - \frac{1}{2}\alpha^2 x^2 - \dots \\ &= 1 - \sum_1^{\infty} b_n x^n; \end{aligned}$$

this series converges for $|x| < 1/2|\alpha|$. If we set $\beta_n = |b_n|/2$, then, since each b is the product of a power of α and a positive numerical coefficient, we see readily from (9) that

$$\begin{aligned} \beta_1 &= |\alpha|/2 \\ \beta_2 &= \beta_1 \beta_1 \\ (10) \quad &\dots \dots \dots \\ \beta_m &= \beta_1 \beta_{m-1} + \beta_2 \beta_{m-2} + \dots + \beta_{m-1} \beta_1 \\ &\dots \dots \dots \end{aligned}$$

and conversely (10) leads to a convergent series.

If we set $|a_n| = \alpha_n$, then from (8)

$$(11) \quad \alpha_1 = |\alpha|, \quad 2\alpha_m \leq \alpha_1 \alpha_{m-1} + \alpha_2 \alpha_{m-2} + \dots + \alpha_{m-1} \alpha_1 + \alpha_{m/2},$$

and, if γ_n is the sequence of positive numbers defined by

$$\gamma_1 = |\alpha|, \quad 2\gamma_m = \gamma_1 \gamma_{m-1} + \gamma_2 \gamma_{m-2} + \cdots + \gamma_{m-2} \gamma_1 + \gamma_{m/2}$$

then

$$(12) \quad \alpha_r \leq \gamma_r, \quad (r = 1, 2, \dots).$$

For, assuming that this inequality holds for $r < m$, as is certainly the case for $m = 2$, then from (11)

$$2\alpha_m \leq \sum_1^{m-1} \alpha_r \alpha_{m-r} + \alpha_{m/2} \leq \sum_1^{m-1} \gamma_r \gamma_{m-r} + \gamma_{m/2} = 2\gamma_m,$$

so that (12) follows by induction. The series (7) for $g(x, \alpha)$ therefore converges if $\sum \gamma_n x^n \equiv g(x, |\alpha|)$ converges.

Let δ be a positive quantity satisfying the conditions

$$\delta > |\alpha|, \quad \delta > 1,$$

and let δ_m be the sequence of increasing positive numbers defined by

$$\delta_1 = \delta, \quad \delta_m = \sum_1^{m-1} \delta_r \delta_{m-r}, \quad (m = 2, 3, \dots),$$

then

$$\gamma_m \leq \delta_m.$$

For this inequality is true for $m = 1$ and, assuming that it is true for $r < m$, we have

$$2\gamma_m = \sum_1^{m-1} \gamma_r \gamma_{m-r} + \gamma_{m/2} \leq \sum_1^{m-1} \delta_r \delta_{m-r} + \delta_{m/2} = \delta_m + \delta_{m/2} < 2\delta_m,$$

since the sequence of δ 's continually increases. Now we have already seen in (10) that $\sum \delta_r x^r$ converges for $|x| < 1/4\delta$; hence $\sum \gamma_n x^n$ converges, and therefore also $\sum \alpha_n x^n$, that is to say, the series (7) converges absolutely for

$$|x| < 1/4|\alpha|, \quad |x| < 1/4.$$

We have therefore proved that there always exists a unique solution of (5) which is regular at the origin.

When α is real and positive, an upper limit for the radius of convergence may be found as follows. The first three terms in the series for $g(x)$ are

$$1 - \alpha x - \frac{\alpha(\alpha+1)}{2} x^2,$$

the remaining terms all having negative coefficients when $\alpha > 0$; hence $g(\alpha^{-1})$ is negative, if the series converges for that value of x . Now $g(0) = 1 > 0$, so that, if the radius of convergence, R , is greater than $1/\alpha$, there is some value ζ of x for which $g(\zeta) = 0$; and, if we put $x = \zeta^{\frac{1}{2}}$ in (5), we have

$$0 = g(\zeta) = 2\alpha\zeta^{\frac{1}{2}} + g^2(\zeta^{\frac{1}{2}}),$$

so that $g(\zeta^{\frac{1}{2}})$ is imaginary. Since this is impossible so long as both ζ and $\zeta^{\frac{1}{2}}$ are inside the circle of convergence, R is certainly less than the smaller of $1/\alpha$ and $1/\alpha^{\frac{1}{2}}$. In particular, if $\alpha > 1$, then $R < 1$. Closer limits are of course obtained by taking more terms of the series. For instance, if $\alpha = 1$ and $x = 2/3$, the sum of the first three terms of the series is negative and therefore the same argument as before shows that $R < 2/3$. Similarly if $\alpha > (\sqrt{13} - 3)/2 = 0.3027 \dots$, the radius R is less than 1.

4. The singularities of $g(x)$. If we write (5) in the form

$$(13) \quad g(x) = 2\alpha x^{\frac{1}{2}} + g^2\left(x^{\frac{1}{2}}\right),$$

we see immediately that, if $x = \zeta$ is a singularity, so is also $x = \zeta^{\frac{1}{2}}$; and similarly all the points $\zeta^{2^{-n}}$ ($n = 1, 2, \dots$) are singularities. If $|\zeta| = 1$, all these points lie on the circle* C_1 , while, if $|\zeta| \neq 1$, the points approach more and more closely to this circle as n increases and at the same time become more and more numerous, since there are 2^n determinations of $\zeta^{2^{-n}}$, and moreover in such a way that every point of C_1 is a limit point of the set of singularities. The circle C_1 therefore forms a natural boundary across which $g(x)$ cannot be continued analytically. Hence *the radius of convergence of the series (7) is never greater than 1 unless it is infinite.*†

In exactly the same way, using (5) in place of (13), it follows that, if ζ is a singularity, so is also ζ^2 unless $g^2(x)$ is regular at $x = \zeta$ which is then a branch point of order 2 at which $g(x) = 0$. If ζ is a singularity for which

* The circle with center at the origin and radius $|x| = r$ will be denoted by C_r or C_r .

† It is shown below that the radius is only infinite when $\alpha = -1$ and $g(x) = 1 + x$.

$|\zeta|$ is a minimum and we assume $|\zeta| < 1$, then $|\zeta| > 0$ and, since $|\zeta^n| < |\zeta|$, it follows that $g^2(x)$ is regular at $x = \zeta$, which is therefore a branch point of order 2 at which $g(x) = 0$. We may therefore set

$$g(x) = (x - \zeta)^{\frac{1}{2}} g_1(x - \zeta),$$

where $g_1(x)$ is regular at $x = \zeta$. Moreover, if $|\zeta|$ is not a minimum but is still less than 1, then, since $|\zeta^n| \rightarrow 0$ as $n \rightarrow \infty$ while $g(x)$ is regular at $x = 0$, $g(x)$ must be regular at ζ^n for some value of n . It then follows from (5) that $g(\zeta)$ is finite and has a finite number of determinations; such singularities are therefore algebraic. We shall now show that, if there is a singularity inside C_1 , there is a *unique* singularity for which $|\zeta|$ is a minimum.

We have already seen in (6) that $h(x) = g(x)/x^{\frac{1}{2}}$ satisfies the equation

$$(14) \quad h(x^2) = 2\alpha + h^2(x).$$

If x_1 and x_2 are two different zeros of $g(x)$ which lie inside C_1 , then

$$(15) \quad h(x_1^{2^n}) = h(x_2^{2^n}),$$

as each is the same polynomial in 2α , e. g., if in (14) $h(x) = 0$, then

$$h(x^2) = 2\alpha,$$

$$h(x^4) = 2\alpha + h^2(x^2) = 2\alpha + 4\alpha^2,$$

and so on. Now $f(x) = 1/h^2(x) = x/g^2(x)$ is regular at $x = 0$, and

$$f'(x) = \frac{g(x) - 2xg'(x)}{g^3(x)},$$

which has the value 1 at $x = 0$. But in (15), $|x_1^{2^n}|$ and $|x_2^{2^n}|$ can be made as small as we please by making n sufficiently large; hence there are an infinity of distinct pairs of points x' and x'' in any neighbourhood of the origin for which

$$\frac{f(x') - f(x'')}{x' - x''} = 0.$$

This is impossible since $f'(0) = 1$; hence $g(x)$ vanishes for at most one value of x within C_1 . We have already shown that $g(x)$, and therefore also $h(x)$,

vanishes at the branch point nearest the origin and hence it follows that there is not more than one such point.

We have therefore shown that $g(x)$ has no singularities within the circle C_1 except possibly branch points of finite order; and, if it has any singularity within C_1 , there is a unique singularity ζ for which $|\zeta|$ is a minimum; this point is a branch point of order 2 at which $g(x) = 0$ and it is the only zero within C_1 . Every point $\zeta^{1/2^n}$ is also a singular point and every singular point within C_1 is of this form.

Exactly the same argument as above may be used to show that $h(x)$ never takes on the same value twice within C_1 .

We have seen above that C_1 in general forms a natural boundary for $g(x)$. There exist, however, solutions which have a simple pole at $x = \infty$ but otherwise behave in the region exterior to C_1 in much the same way as $g(x)$ does in the interior region, or, more explicitly, if ζ_1, ζ_2, \dots are the singularities of $g(x)$ within C_1 , then

$$\bar{g}(x) = xg\left(\frac{1}{x}\right)$$

is a solution of (5) whose only singularities outside C_1 are $\zeta_1^{-1}, \zeta_2^{-1}, \dots$ and a simple pole at $x = \infty$. For, if $|x| > |\zeta^{-1}|$, ζ being as before the singularity of $g(x)$ whose modulus is least, then

$$\begin{aligned}\bar{g}(x^2) &= x^2 g\left(\frac{1}{x^2}\right) = x^2 \left[\frac{2\alpha}{x} + g^2\left(\frac{1}{x}\right) \right] \\ &= 2\alpha x + \left[xg\left(\frac{1}{x}\right) \right]^2 = 2\alpha x + \bar{g}^2(x).\end{aligned}$$

In particular, when $g(x) = 1 + x$, we have $\bar{g}(x) \equiv g(x)$.

It is readily seen that no entire function except $1 + x$ can be a solution of (5). For differentiating the terms of this equation we have

$$\begin{aligned}g(x^2) &= 2\alpha x + [g(x)]^2 \\ 2xg'(x^2) &= 2\alpha + 2gg', \\ 4x^2g''(x^2) + 2g'(x^2) &= 2gg'' + 2g'^2 \\ &\dots \dots \dots \\ 2^n x^n g^{(n)}(x^2) + \frac{n(n-1)}{2} 2^{n-1} x^{n-2} g^{(n-1)}(x^2) + \dots \\ &= 2g(x)g^{(n)}(x) + 2ng'(x)g^{(n-1)}(x) + \dots\end{aligned}$$

Hence the value of $g^{(n)}(1)$ is determined uniquely in terms of $g(1)$ unless $g(1) = 2^{n-1}$, i. e., $2\alpha = 2^{n-1} - 2^{2n-2}$ for any positive integral value of n . For these values of α , we have, on putting $x = -1$ in (5),

$$g(1) = -2^{n-1} + 2^{2n-2} + [g(-1)]^2$$

or

$$[g(-1)]^2 = 2^{n-1} + 2^{n-1} - 2^{2n-2} = -(2^{2n-2} - 2^n) < 0$$

unless $n = 1, 2$. Since the series for $g(x)$ has real values when α and x are real, $g(-1)$ cannot have a negative square so that we need only consider the cases $n = 1, 2$. For $n = 2$, we have $\alpha = -1$, and, as we have already seen, $g(x) = 1 + x$ which is an entire function; for $n = 1$, $\alpha = 0$ and $g(x) = 1$, again an entire function; we may therefore exclude these two trivial cases.

Now $\bar{g}(x)$ satisfies the same functional equation as $g(x)$ and $\bar{g}(1) = g(1)$, and therefore the derivatives of $\bar{g}(x)$ have the same values at $x = 1$ as those of $g(x)$. But, if $g(x)$ is an entire function, $\bar{g}(x)$ is regular at $x = 1$; it follows therefore that the series defining $g(x)$ and $\bar{g}(x)$ are identical. This is impossible as $g(x)$ is regular at $x = 0$ while $\bar{g}(x)$ has a singularity there except for the cases $\alpha = 0, 1$ considered above. We have therefore shown that $g(x)$ is not an entire function except when $\alpha = 0$, $g(x) = 1$, and $\alpha = -1$, $g(x) = 1 + x$.

5. The radius of convergence. The radius of convergence of (7) can be calculated by means of (14) under certain restrictions and, when α is real, also the value of x which corresponds to a given real value of $g(x)$.

By successive applications of (14) we have

$$\begin{aligned} h(x^2) &= 2\alpha + h^2(x) \\ h(x^4) &= 2\alpha + (2\alpha + h^2(x))^2 \\ &\dots \dots \dots \\ h(x^{2^n}) &= 2\alpha + (2\alpha + \dots + (2\alpha + h^2(x))^2 \dots)^2, \end{aligned}$$

where in $h(x^{2^n}) = g(x^{2^n})/x^{2^{n-1}}$ the term 2α occurs n times. If we set

$$(17) \quad k_n(x) = |h(x^{2^n})|^{\frac{1}{2^{n-1}}} = \frac{|g(x^{2^n})|^{\frac{1}{2^{n-1}}}}{|x|},$$

then, since, if $|x| < 1$, $x^{2^n} \rightarrow 0$ as $n \rightarrow \infty$ and $g(0) = 1$, it follows that

$$(17') \quad \lim_{n \rightarrow \infty} k_n(x) = \frac{1}{|x|}.$$

If we replace $h(x)$ by λ in (16) and write*

$$(18) \quad \begin{aligned} h_0 &= \lambda, & k_0 &= |\lambda^2|, \\ h_1 &= 2\alpha + \lambda^2, & k_1 &= |2\alpha + \lambda^2|, \\ &\dots & &\dots \\ h_n &= 2\alpha + h_{n-1}^2, & k_n &= |h_n|^{\frac{1}{2^{n-1}}}, \end{aligned}$$

then k_n approaches a definite limit as $n \rightarrow \infty$ provided λ is admissible as a value of $h(x)$ for $|x| < 1$. The convergence of k_n to its limit is, as a rule, rapid for real values of α which are not too small, and this furnishes a practical method of calculating the zero $x = \zeta$, which corresponds to $\lambda = 0$, provided we are able otherwise to determine that it lies inside C_1 .

The values of ζ calculated in this manner for certain values of α are

α	ζ	α	ζ
0.25	0.9292	5.00	0.09533
0.50	0.6654	10.00	0.04880
1.00	0.4027	50.00	0.00995
2.00	0.2230		

The convergence of k_n to its limit can also be determined independently under certain conditions. We shall suppose in the first case that for some value of r

$$(19) \quad |h_r| \geq 1 + |2\alpha|^{\frac{1}{2}}$$

so that

$$|h_r|^2 \geq 1 + 2|2\alpha|^{\frac{1}{2}} + |2\alpha| > |2\alpha|;$$

then

$$|h_{r+1}| = |2\alpha + h_r^2| > |h_r|^2 - |2\alpha| \geq 1 + 2|2\alpha|^{\frac{1}{2}},$$

* When it is necessary to indicate the values of λ and 2α explicitly, we shall write $h_n(\lambda, 2\alpha)$ in place of h_n .

whence it follows easily by induction that

$$(20) \quad |h_{r+n}| \geq 1 + 2^n |2\alpha|^{\frac{1}{2}};$$

and therefore if n is sufficiently large,* $|h_n| > |2\alpha|$.

Now

$$\begin{aligned} \frac{k_n}{k_{n-1}} &= \left| \frac{h_n}{h_{n-1}^2} \right|^{\frac{1}{2^{n-1}}} = \left| 1 + \frac{2\alpha}{h_{n-1}^2} \right|^{\frac{1}{2^{n-1}}} \leq \left(1 + \frac{|2\alpha|}{|h_{n-1}^2|} \right)^{\frac{1}{2^{n-1}}} \\ &\leq (1 + |2\alpha|)^{\frac{1}{2^{n-1}}} < 1 + \frac{|2\alpha|}{2^{n-1}} \end{aligned}$$

for n sufficiently large. The infinite product $\prod k_n/k_{n-1}$ therefore converges and so k_n approaches a definite, finite limit.

If $|2\alpha| \geq 2.25$, it is easily shown that (19) is satisfied for $\lambda = 0$ and $r = 2$.

The convergence follows in the same way if it is known that $|h_n| \geq \varepsilon > 0$ for all values of $n > r$, ε being independent of n .

If $\lambda = 0$ is a possible value of λ and z , $|z| < 1$, the corresponding value of x , then as before

$$h(z^2) = 2\alpha,$$

$$h(z^4) = 2\alpha + (2\alpha)^2$$

$$\dots \dots \dots$$

$$h(z^{2^n}) = 2\alpha + h^2(z^{2^{n-1}}).$$

Hence $h(z^{2^n})$ is a polynomial in α of degree 2^{n-1} whose coefficients are positive integers which do not depend on z or α ; this polynomial we shall denote by $p_n(2\alpha)$. If $\mu = 2\alpha$ is a root of $p_n(\mu) = 0$, then $h(z^{2^n}) = 0$, so that z^{2^n} is a root of $g(x)$. Hence $(z^{2^n})^{2^n}$ is also a root, and so on. This, however, is impossible if $|z| < 1$, as the origin would then be an essential singularity of $g(x)$; we can therefore conclude that, *if 2α is a root of any of the polynomials $p_n(\mu)$, $g(x)$ vanishes at no point within C_1 which is therefore the circle of convergence of the series (7), except in the two trivial cases in which this series represents an entire function.*

6. The polynomials $h_n(\lambda, \mu)$ and $p_n(\mu)$. The polynomial $h_n(\lambda, \mu)$ is the polynomial defined in (18) with μ in place of 2α , i. e.,

$$(21) \quad h_0(\lambda, \mu) = \lambda, \quad h_n(\lambda, \mu) = \mu + h_{n-1}^2(\lambda, \mu),$$

* If $|h_n| > 1 + |2\alpha|^{\frac{1}{2}}$ for every n greater than a certain value, then evidently $\lim k_n \leq 1$.

while $p_n(\mu) = h_n(0, \mu)$ so that

$$(21') \quad p_1(\mu) = \mu, \quad p_n(\mu) = \mu + p_{n-1}^2(\mu).$$

The following properties follow readily from these definitions.

In the first place we have

$$(22) \quad h_n(\lambda, \mu) = h_{n-r}(h_r(\lambda, \mu), \mu) = p_{n-r}(\mu) + q_{nr}(\lambda, \mu) h_r^2(\lambda, \mu),$$

where $q_{nr}(\lambda, \mu)$ is a polynomial in λ, μ with positive integral coefficients; hence in particular

$$(22') \quad p_n(\mu) = p_{n-r}(\mu) + q_{nr}(0, \mu) p_r^2(\mu).$$

From these equations we deduce immediately that

(i) $h_n(\lambda, \mu)$ and $h_r(\lambda, \mu)$ have no common factor; and if s is the H. C. F. of n and r , $p_s(\mu)$ is the H. C. F. of $p_n(\mu)$ and $p_r(\mu)$.

To prove the first part of this lemma we need only observe that from (22) every common factor of h_n and h_r is a factor of $p_{n-r}(\mu)$ and therefore does not contain λ . This is, however, impossible since the coefficient of the highest power of λ in h_r is unity.

To prove the second part we observe from (22') that every common factor of p_n and p_r is a factor of p_{n-r} ; and hence, by a repetition of this argument, it is a factor of p_s . It only remains, therefore, to prove that p_s is a factor of p_{ks} , k being any integer. Putting $n = 2s, 3s, \dots, r = s, 2s, \dots$ in (22'), we have

$$p_{2s} = p_s + q_{2s,s} p_s^2, \quad p_{3s} = p_{2s} + q_{3s,s} p_s^2,$$

and so on; from which the required result follows immediately. This also shows that $p_1 = \mu$ is a factor of every $p_n(\mu)$ as is of course obvious otherwise.

(ii)

$$(23) \quad h_n(\lambda, \mu) - h_r(\lambda, \mu) = (h_{n-1} + h_{r-1})(h_{n-2} + h_{r-2}) \cdots (h_{n-r} + h_0)(h_{n-r} - h_0),$$

$$(23') \quad p_n(\mu) - p_r(\mu) = (p_{n-1} + p_{r-1})(p_{n-2} + p_{r-2}) \cdots (p_{n-r+1} + p_1) p_{n-r}^2.$$

For

$$\begin{aligned} h_s(\lambda, \mu) - h_t(\lambda, \mu) &= h_{s-1}^2(\lambda, \mu) - h_{t-1}^2(\lambda, \mu) \\ &= [h_{s-1}(\lambda, \mu) + h_{t-1}(\lambda, \mu)] [h_{s-1}(\lambda, \mu) - h_{t-1}(\lambda, \mu)]. \end{aligned}$$

An immediate consequence of (23) is that $h_n(\lambda, \mu)$, ($n = k, k+1, k+2, \dots$) all have the same value for values of λ and μ for which

$$h_r(\lambda, \mu) + h_{r-1}(\lambda, \mu) = 0, \quad (r = 1, 2, \dots k),$$

an important particular case of which is

$$p_n(-2) = 2, \quad (n \geq 2).$$

(iii) Differentiating (21) with regard to λ^2 and μ , we readily prove by induction that

$$(24) \quad \frac{\partial h_n}{\partial \mu} = 1 + 2h_{n-1} + 2^2 h_{n-1} h_{n-2} + 2^3 h_{n-1} h_{n-2} h_{n-3} + \dots \\ \dots + 2^{n-1} h_{n-1} h_{n-2} \dots h_1,$$

$$(24') \quad \frac{\partial h_n}{\partial \lambda^2} = 2^{n-1} h_{n-1} h_{n-2} \dots h_1,$$

an interesting particular case of which is $p'_n(-2) = -(2^{2n-1} + 1)/3$, ($n \geq 2$).

(iv) If

$$(25) \quad |\lambda^2 + \mu| \geq \frac{1}{2} + \sqrt{|\mu| + \frac{1}{4}},$$

then

$$(26) \quad |h_n(\lambda, \mu)| \geq |\lambda^2 + \mu|^2 - |\mu| \geq |\mu|^{\frac{1}{2}}, \quad (n \geq 2),$$

and, in particular, if $|\mu| \geq 2$,

$$(26') \quad |p_n(\mu)| \geq |\mu|^2 - |\mu| \geq |\mu|^{\frac{1}{2}}, \quad (n \geq 2).$$

If (25) is satisfied, then

$$|\mu| + |\mu|^{1/2} < |\mu| + \frac{1}{2} + \sqrt{|\mu| + \frac{1}{4}} = (\frac{1}{2} + \sqrt{|\mu| + \frac{1}{4}})^2 \geq 1;$$

hence

$$|\lambda^2 + \mu|^2 \geq |\mu| + |\mu|^{1/2},$$

or

$$|\lambda^2 + \mu|^2 - |\mu| \geq |\mu|^{1/2} \geq 0.$$

Now, for $n = 2$,

$$|h_2(\lambda, \mu)| = |\mu + (\lambda^2 + \mu)^2| \geq |\lambda^2 + \mu|^2 - |\mu| \geq |\mu|^{1/2}.$$

Let us assume, therefore, that (26) is true for 2, 3, ... n ; then

$$\begin{aligned} |h_{n+1}(\lambda, \mu)| &= |h_n^2(\lambda, \mu) + \mu| \geq |h_n^2(\lambda, \mu)| - |\mu| \\ &\geq |\lambda^2 + \mu|^4 - 2|\mu| |\lambda^2 + \mu|^2 + |\mu|^2 - |\mu| \\ &= |\lambda^2 + \mu|^2 - |\mu| + [|\lambda^2 + \mu|^2 - (|\mu| + \tfrac{1}{2})]^2 - |\mu| - \tfrac{1}{4} \\ &\geq |\lambda^2 + \mu|^2 - |\mu| \geq |\mu|^{1/2} \end{aligned}$$

by (25). Equation (26) then follows by induction.

If $\lambda = 0$, (25) becomes

$$|\mu| \geq \tfrac{1}{2} + \sqrt{|\mu| + \tfrac{1}{4}},$$

or $|\mu|^2 \geq 2|\mu|$, i. e., $|\mu| \geq 2$; and (26) becomes (26').

Since $|\lambda^2 + \mu| \geq ||\lambda|^2 - |\mu||$, (25) is satisfied by

$$|\lambda|^2 \geq |\mu| + \tfrac{1}{2} + \sqrt{|\mu| + \tfrac{1}{4}}, \quad |\lambda|^2 > |\mu|,$$

or

$$|\lambda|^2 \leq |\mu| - \tfrac{1}{2} - \sqrt{|\mu| + \tfrac{1}{4}}, \quad |\lambda|^2 < |\mu| \leq 2.$$

(v) It follows immediately from (26') that *the absolute value of every root of $p_n(\mu)$, except $\mu = 0$, is less than 2 and therefore corresponds to a value of $|\alpha| < 1$ and, if α is real, to a value of α between -1 and 0 .*

(vi) *If μ_n is the real negative root of p_n of greatest absolute value, there is one, and only one, real root μ_{n+1} of p_{n+1} between -2 and μ_n .*

When $\mu = -2$, we have already seen that $p_n = 2 > 0$, and if $\mu = \mu_{n-1}$, then $p_n = \mu_{n-1} + p_{n-1}^2 = \mu_{n-1} < 0$; there is therefore at least one real root of p_n between -2 and μ_{n-1} .

If we differentiate (21') twice, we get

$$p_n''(\mu) = 2[p_{n-1}(\mu)]^2 + 2p_{n-1}(\mu)p_{n-1}''(\mu),$$

so that, if μ is real, p_n'' is positive if p_{n-1} and p_{n-1}'' have the same sign. For $n = 3$, $\mu_3 = -1$; and, for $\mu < -1$, we evidently have $p_3'' \geq 0$, so that to the left of $\mu = \mu_3$ both p_3 and p_3'' are positive. It follows that, to the left of μ_3 , $p_4'' \geq 0$; and so on. We can therefore conclude that p_n has one, and only one, real root between -2 and μ_{n-1} as otherwise p_n'' , which is equal to or greater than 0, would change sign for some value of μ between these limits.

It is also easy to show that $\mu_n \rightarrow -2$ as $n \rightarrow \infty$. For

$$p_n(-2) = 2, \quad p_n(\mu_{n-1}) = \mu_{n-1} < -1,$$

while $p_n''(\mu) \geq 0$ between -2 and μ_{n-1} so that the graph of $p_n(\mu)$ is concave upwards between these limits; μ_n therefore lies to the left of the line joining $(-2, 2)$ to $(\mu_{n-1}, -1)$, whence

$$\mu_{n-1} - \mu_n > \frac{1}{3}(\mu_{n-1} + 2).$$

(vii) If μ and λ are real and positive, and if $\mu \leq \frac{1}{4}$ and $\mu + \lambda^2 - \lambda < 0$ or $\lambda < \mu + \lambda^2 < \frac{1}{4}$, then

$$\lim_{n \rightarrow \infty} h_n(\lambda, \mu) = \frac{1}{2} - \sqrt{\frac{1}{4} - \mu}.$$

For all other positive values of μ and λ , $h_n(\lambda, \mu) \rightarrow \infty$ as $n \rightarrow \infty$.

From (23) with $r = n - 1$ we have

$$h_n - h_{n-1} = (h_{n-1} + h_{n-2})(h_{n-2} + h_{n-3}) \cdots (h_1 + h_0)(\mu + \lambda^2 - \lambda).$$

If $\mu + \lambda^2 - \lambda < 0$, which requires $\mu < 1/4$, the h 's therefore form a decreasing sequence of positive quantities and so approach a finite limit. If $\mu + \lambda^2 - \lambda > 0$, they form an increasing sequence. If the sequence of h 's has a finite limit, then, since $h_n = \mu + h_{n-1}^2$, we must have $l = \mu + l^2$. Moreover, since $h_{n-1} + h_{n-2} \rightarrow 2l$, we see immediately from (23) that $l < 1/2$, and hence $l = \frac{1}{2} - \sqrt{\frac{1}{4} - \mu}$. A finite limit can therefore only exist if $\mu < 1/4$.

If $\mu < 1/4$ and $h_1 \equiv \mu + \lambda^2 + 1/2$, then

$$h_2 = \mu + (\mu + \lambda^2)^2 < \frac{1}{2},$$

and therefore, by an easy induction, $h_n < 1/2$ for every n . On the other hand, if $\lambda < \mu + \lambda^2 > 1/2$, the h 's form an increasing sequence which cannot have a finite limit since $h_{n-1} + h_{n-2}$ is greater than unity.

(viii) If $\mu > 1/4$, $\lambda > 0$ and $\frac{1}{x} = \lim_{n \rightarrow \infty} k_n(\lambda, \mu)$, then $\lambda = h(x, \mu)$.

In the first place, $k(\lambda, \mu) \equiv \lim_{n \rightarrow \infty} k_n(\lambda, \mu)$ exists and is finite. For, under the given conditions, h_n increases indefinitely with n and hence the condition of (19) and (20) of § 5 that $h_n > 1 + 2^{n-r} \mu^{\frac{1}{2}}$ is satisfied for n greater than some finite value of r . The value of the limit is moreover greater than unity; for we can write

$$h_n > (1 + \epsilon)^{2n-1}, \quad (\epsilon > 0)$$

for some n sufficiently large and ϵ sufficiently small, whence

$$h_{n+1} > \mu + (1 + \epsilon)^{2n} > (1 + \epsilon)^{2n},$$

so that the inequality also holds, with the same ϵ , for every subsequent value of n .

We shall now show that $\partial k_n / \partial \lambda$ approaches a finite value as $n \rightarrow \infty$. From (17) and (24') we have

$$\frac{1}{2\lambda} \frac{\partial k_n}{\partial \lambda} = \frac{\partial k_n}{\partial \lambda^2} = k_n \frac{h_{n-1} h_{n-2} \cdots h_1}{h_n} \equiv k_n(\lambda) q_n(\lambda)$$

say. Now

$$\frac{q_{n+1}}{q_n} = \frac{h_n^2}{h_{n+1}} = \frac{1}{1 + \frac{\mu}{h_n^2}},$$

and $\sum \mu / h_n^2$ converges since $h_n > 1 + 2^{n-r} \mu^{\frac{1}{2}}$ for n sufficiently large. Hence q_n approaches a definite finite limit as $n \rightarrow \infty$, and this limit is never zero. Moreover, k_n and q_n approach their limits uniformly as regards λ since in both cases the convergence was obtained by comparison with series which are independent of λ . Hence

$$\frac{\partial k(\lambda, \mu)}{\partial \lambda} = 2\lambda k(\lambda, \mu) q(\lambda, \mu).$$

It follows that, in any interval $0 < \epsilon \leq \lambda \leq N$ in which ϵ is as small and N as great as we please, $x \equiv 1/k(\lambda, \mu)$ is a uniformly continuous function of λ and

possesses a derivative which is nowhere zero. There exists therefore a unique single-valued inverse function $\lambda = H(x, \mu)$. Now we saw in § 5 that $k(\lambda, \mu) = 1/x$ if λ is admissible as a value of $h(x, \mu)$; also $h(x, \mu) \rightarrow \infty$ as $x \rightarrow 0$; hence, if N is taken large enough, the range of values for $h(x, \mu)$ will overlap that for $H(x, \mu)$, and in the common part of their ranges these two functions have the same value. Both functions satisfy the same functional equation, namely $h(x^2) = \mu + h^2(x)$; hence, if a is so small that a and $a^{1/2}$ both lie in the common range, so will also $a^{1/4}$ provided always that $h(x, \mu)$ and $H(x, \mu)$ do not vanish; and so on. The two functions therefore have the same range of definition and this range is from ϵ to the point at which they vanish or, if they do not vanish, from ϵ to 1. We can then use (17') to calculate the zero of $g(x)$ provided $\mu > 1/4$, and the radius of convergence of $g(x)$ is less than unity for real values of $2\alpha \equiv \mu > 1/4$, and it is equal to unity for $0 < \mu \leq 1/4$.

7. Solutions which have a singular point at the origin. If $g(x)$ has a pole of order n at $x = 0$, then

$$G(x) = x^n g(x)$$

is regular there and $G(0) \neq 0$. The function $G(x)$ satisfies the functional equation

$$(27) \quad G(x^2) = 2\alpha x^{2n+1} + G^2(x),$$

so that $G(0) = 1$. Since $2n + 1$ is integral whenever n is an integral multiple of $1/2$, the discussion of this equation will also embrace those solutions which have an algebroid pole or zero whose order is of the form $m/2$.

In place of (27) we shall consider the more general equation

$$(28) \quad F(x^2) = 2(\alpha^0 + \alpha'x + \alpha''x^2 + \dots + \alpha^{(n)}x^n) + F^2(x),$$

and we shall only consider solutions which are regular at the $x = 0$. Substituting in this equation

$$(29) \quad F(x) = a_0 - a_1x - a_2x^2 - a_3x^3 - \dots$$

so that it also holds for every value of m . The proof that the series converges is then exactly as in § 3.

(ii) Suppose that $1/\alpha_0 > 1$. Then, γ being chosen as before, let

$$\gamma_1 = \gamma, \quad 2\gamma_m = \sum \gamma_r \gamma_{m-r} + \frac{1}{\alpha_0} \gamma_{m/2},$$

so that

$$2\beta_m \leq \sum \gamma_r \gamma_{m-r} + \frac{1}{\alpha_0} \gamma_{m/2} = 2\gamma_m, \quad (m > n).$$

Let δ be a positive quantity satisfying the conditions

$$\delta \geq \alpha_0 \gamma, \quad \delta \geq 1,$$

and set $\delta_1 = \delta$, $\delta_m = \sum \delta_r \delta_{m-r}$; then $\delta_m \geq \alpha_0^m \gamma_m$. This inequality is true when $m = 1$; suppose it is true for all values of the subscript up to $m-1$, then

$$\begin{aligned} 2\gamma_m &= \sum \gamma_r \gamma_{m-r} + \frac{1}{\alpha_0} \gamma_{m/2} \leq \sum \left(\frac{1}{\alpha_0}\right)^m \delta_r \delta_{m-r} + \left(\frac{1}{\alpha_0}\right)^{\frac{m+2}{2}} \delta_{m/2} \\ &= \left(\frac{1}{\alpha_0}\right)^m \delta_m + \left(\frac{1}{\alpha_0}\right)^{\frac{m+2}{2}} \delta_{m/2}. \end{aligned}$$

But $\delta_{m/2} \leq \delta_m$ and $m > (m+2)/2$; hence the inequality holds for every value of m . It then follows exactly as before that (29) converges for $|x| < \alpha_0/4\delta$.

It is not difficult to show* that no solution of (29) can be an entire function unless it is a polynomial. It then follows that the original functional equation (5) cannot have a solution of the form

$$g(x) = a_m x^m + \dots + a_0 + \sum_1^{\infty} a_{-r} x^{-r}$$

in which the infinite series does not terminate. For, if there were such a solution, $G\left(\frac{1}{x}\right) = g(x)/x^m$ would be an entire function of $z = 1/x$ which

* One method of proof is to consider the manner in which $|F(x)|$ increases with $|x|$.

satisfies the equation

$$G(z^2) = 2\alpha z^{2m-1} + G^2(z)$$

and is not a polynomial.

8. **Conclusion.** The method of calculating the zeros of $g(x)$ which was given in § 5 may also be used to calculate the value of $g(x)$ for any real x and α . When this is carried out, it is immediately seen that comparatively few steps of the limiting process are required to give a fairly accurate result. If the number of steps required for a given degree of accuracy has been ascertained, the process may be profitably inverted, e. g., if four steps are sufficient, we may set

$$h(x) = [([x^{-8} - 2\alpha]^{\frac{1}{2}} - 2\alpha)^{\frac{1}{2}} - 2\alpha]^{\frac{1}{2}} - 2\alpha]^{\frac{1}{2}},$$

or in general

$$h(x) = [[\dots [x^{-2^{n-1}} - 2\alpha]^{\frac{1}{2}} - 2\alpha]^{\frac{1}{2}} \dots - 2\alpha]^{\frac{1}{2}} - 2\alpha]^{\frac{1}{2}},$$

where the square root is extracted n times. If n is taken so large that $[x^{-2^{n-1}} - 2\alpha]^{\frac{1}{2}} = x^{-2^{n-2}}$ to the degree of accuracy required, then nothing is gained by increasing the number of steps beyond n .

PRINCETON, N. J.,
December, 1921.